

Cluster algebras of type C via cluster tubes ¹

Yu Zhou and Bin Zhu

Department of Mathematical Sciences
Tsinghua University
100084 Beijing, P. R. China
E-mail: yu-zhou06@mails.tsinghua.edu.cn

Department of Mathematical Sciences
Tsinghua University
100084 Beijing, P. R. China
E-mail: bzhu@math.tsinghua.edu.cn

Abstract

Cluster tubes are 2–Calabi–Yau triangulated categories which contain no cluster tilting objects, but maximal rigid objects. Fix a maximal rigid object T in the cluster tube C_n of rank n . For any indecomposable rigid object M in C_n , we give an analogous X_M of Caldero–Chapton’s formula (or Palu’s cluster character formula) by using the geometric information of M . We show that $X_M, X_{M'}$ satisfy the mutation formula when M, M' form an exchange pair, and that $X_\gamma : M \mapsto X_M$ gives a bijection from the set of indecomposable rigid objects in C_n to the set of cluster variables of cluster algebra of type C_{n-1} , which induces a bijection between the set of (basic) maximal rigid objects in C_n and the set of clusters. This strengthens a surprising result proved recently by Buan–Marsh–Vatne that the combinatorics of maximal rigid objects in the cluster tube C_n encode the combinatorics of the cluster algebra of type B_{n-1} since the combinatorics of cluster algebras of type B_{n-1} or of type C_{n-1} are the same by one of results of Fomin–Zelevinsky in [FZ2].

Key words. Cluster tubes; Maximal rigid objects; Cluster variables; cluster algebras of type C .

Mathematics Subject Classification. 16G20, 16G70.

1 Introduction

Cluster algebras were introduced around 2001 by Fomin–Zelevinsky [FZ1] in order to give an algebraic and combinatorial framework for the canonical basis of quantum groups and for the notion of total positivity for semisimple algebraic groups, see [FZ4] for a nice survey on this topic and its background. Since they introduced, interesting connections between such algebras and several branches of mathematics have emerged. In the categorification theory of cluster algebras, cluster categories [BMRRT, CCS, K1, Am, Du], (stable) module categories over preprojective algebras [GLS1, GLS2, BIRS] play a central role. They all have cluster tilting objects, and these cluster tilting objects model the clusters of the corresponding cluster algebras via Caldero–Chapton’s formula [CC] in the case of cluster categories or Geiß–Leclerc–Shröer’s map [GLS1] in the case of preprojective algebras. This motivates the study of arbitrary 2–Calabi–Yau triangulated categories with cluster tilting objects (subcategories). Palu defined the cluster character for the 2–CY triangulated categories which have cluster tilting objects and all cluster tilting objects have no loops or 2–cycles [Pa1] (see also [FK]). It was proved in [IY] [BIRS] that one can mutate cluster tilting object T (respectively maximal rigid object) at any indecomposable direct summand to get a new cluster tilting object $\mu_i(T)$ (resp. maximal rigid object) via exchange triangles in a 2–CY triangulated category C . For any maximal rigid object T , one can associate an integer matrix A_T by using the exchange triangles (where A_T we define is the transpose of

¹Supported by the NSF of China (Grants 10771112)

B_T defined in [BMV]). If A_T and $A_{\mu_i(T)}$ are related by Fomin-Zelevinsky's matrix mutation for any maximal rigid object T , then we call that the maximal rigid objects form a cluster structure in C [BMV, BIRS]. In [BMV], Buan-Marsh-Vatne showed that maximal rigid objects without 2-cycles form a cluster structure in C (see also [BIRS]). Cluster tilting objects and maximal rigid objects are important objects in 2-CY triangulated categories. They have many nice properties, see for example [I], [KR], [KZ], [IY], [DK], [V], [Y], [ZZ]. Cluster tilting objects are obvious maximal rigid, but the converse is not true, see [BIKR] for the first examples. Cluster tube provide the second examples, in which maximal rigid objects contains loops, contains no 2-cycles [BMV]. Cluster tubes, denoted by C_n , is by definition, the orbit category by $\tau^{-1}[1]$ of derived category of the hereditary abelian category of nilpotent representations of a quiver with underlying graph $\tilde{A}_{n-1}$ and with cyclic orientation. It is a 2-CY triangulated category [BKL1-2]. In [BMV], a classification of maximal rigid objects in the cluster tube C_n is given. The maximal rigid objects is proved to form a cluster structure. Furthermore they use the geometric description of the exchange graph of cluster algebras of type B_{n-1} in [FZ2] to prove that there is a bijection between the set of indecomposable rigid objects in cluster tube C_n and the set of cluster variables of cluster algebra of type B_{n-1} . Under this bijection, maximal rigid objects go to clusters [BMV]. Since the cluster combinatorics of cluster algebras of type C_{n-1} is the same as that of cluster algebras of type B_{n-1} by Proposition 3.15 in [FZ2], there is also a bijection between the set of indecomposable rigid objects in cluster tube C_n and the set of cluster variables of cluster algebra of type C_{n-1} .

The aim of the paper is give a analogue of Caldero-Chapoton's formula [CC] (or Palu's character [Pa1]) for cluster tubes. Fix a maximal rigid object T in C_n with the matrix A_T associated with T being a skew-symmetrizable matrix of type C_{n-1} . For any indecomposable rigid object M in C_n , with respect to T , we define an analogue of the Caldero-Chapoton's formula X_M defined for cluster categories in [CC], see also [Pa1]. We prove that the formula X_M satisfies mutation formula: i.e. if M and M^* are indecomposable rigid objects such that $M \oplus N$ and $M^* \oplus N$ are maximal rigid objects in C_n , then $X_M \cdot X_{M^*} = X_E + X_{E'}$ where E, E' are the middles of the exchange triangles: $M \rightarrow E \rightarrow M^* \rightarrow M[1]$, $M^* \rightarrow E' \rightarrow M \rightarrow M^*[1]$. We note here that the dimension of $\text{Ext}^1(M, M^*)$ maybe 2 (compare to the cases considered before in [CC],[Pa1],[FK],[Pla1-2], the k -dimension of $\text{Ext}^1(M, M^*)$ is always one). Thus X_γ gives a bijection from the set of indecomposable rigid objects in C_n to the set of cluster variables of cluster algebras of type C_{n-1} . This gives an explicit bijection parallel as that given by Buan-Marsh-Vatne [BMV] for type B_{n-1} (since there is a nature bijection between type B_{n-1} and type C_{n-1} , see [FZ2]). The algebra generated by the X_M , where M runs over indecomposable rigid objects in C_n is isomorphic to the cluster algebras of type C_{n-1} . Ming Ding and Xu Fan also defined an analogue map for cluster tubes and gave multiplication formulas but it can not be used to realize the cluster algebra of type C or B [DX].

The paper is organized as follows: In Section 2, we recall some basics on cluster algebras and 2-CY triangulated categories. In particular, we recall the definition of cluster tubes and basic description on indecomposable rigid objects in cluster tubes [BMV]. In Section 3, fix a maximal rigid object T in C_n , We calculate the index of any indecomposable rigid object M respect to T (defined in [Pa1, DK, Pla]) and define the analogue X_M of CC-maps or Palu's map for indecomposable rigid object M with respect to T . This map X_γ is called cluster map. Using the structure of the cluster tube C_n , we divide the set of indecomposable rigid objects into three disjoint subsets. Using the structure of endomorphism algebras of T , we calculate the explicit formula of

X_M according to which subset M belongs to. In Section 4, we prove that X_M, X_{M^*} satisfy the mutation formula when M, M^* form a mutation pair. Using the mutation triangles, we explain the matrix A_T associated with T is a skew-symmetrizable matrix of type C_{n-1} . We prove that the map X_γ gives a bijection between the set of indecomposable rigid objects in C_n and set of cluster variables of A_T , which also induces a bijection between the set of maximal rigid objects and the set of clusters of A_T . It follows that the cluster algebra generated by X_M , where M runs over all indecomposable rigid objects, is isomorphic to the cluster algebra of type C_{n-1} .

2 Preliminaries

We recall some basic notation on cluster algebras which can be found in the papers by Fomin and Zelevinsky [FZ1-3]. The cluster algebras we deal with in this paper are without coefficients.

Let $\mathcal{F} = \mathbf{Q}(u_1, u_2, \dots, u_n)$ be the field of rational functions in indeterminacies $u_1, u_2, \dots, u_n$. Set $\underline{u} = \{u_1, u_2, \dots, u_n\}$. Let $A = (a_{ij})$ be an $n \times n$ skew-symmetrizable integer matrix. For any $k \in [1, n]$, where $[1, n]$ denotes the set of the numbers $1, 2, \dots, n$, the mutation $\mu_k(A)$ of A in direction k is by definition, an integer matrix $A' = (a'_{ij})$,

$$a'_{ij} = \begin{cases} -a_{ij} & \text{if } i = k \text{ or } j = k, \\ a_{ij} + \frac{|a_{ik}||a_{kj} + a_{ik}||a_{kj}|}{2} & \text{otherwise.} \end{cases}$$

It is also a skew-symmetrizable matrix.

A pair $(\underline{x}, A)$, where $\underline{x} = \{x_1, x_2, \dots, x_n\}$ is a transcendence base of $\mathcal{F}$ and A is an $n \times n$ skew-symmetrizable integer matrix, is called a seed.

A mutation $\mu_k(\underline{x}, A)$ of a seed $(\underline{x}, A)$ in direction k is a new seed $(\underline{x}', A')$, where $A' = \mu_k(A)$, and $\underline{x}' = \underline{x} - \{x_k\} \cup \{x'_k\}$, x'_k is defined in the following mutation formula:

$$x_k x'_k = \prod_{a_{ik} > 0} x_i^{a_{ik}} + \prod_{a_{ik} < 0} x_i^{-a_{ik}}.$$

The "mutation equivalence $\approx$ " is an equivalent relation on the set of all seeds generated by the mutation.

The cluster algebra $\mathcal{A}_A$ associated to the skew-symmetrizable matrix A is by definition the subalgebra of $\mathcal{F}$ generated by all x_i in $\underline{x}$ such that $(\underline{x}, A') \approx (\underline{u}, A)$. Such $\underline{x} = (x_1, x_2, \dots, x_n)$ is called a cluster of the cluster algebra $\mathcal{A}_A$ or simply of the matrix A , and any x_i is called a cluster variable. The set of all cluster variables is denoted by χ_A . If the set χ_A is finite, then the cluster algebra $\mathcal{A}_A$ is called of finite type. For any skew-symmetrizable integer matrix A , one can define the Cartan part C_A of A as follow: $C_A = (c_{ij})_{n \times n}$, where

$$c_{ij} = \begin{cases} -|a_{ij}| & \text{if } i \neq j, \\ 2 & i=j. \end{cases}$$

It was proved by Fomin-Zelevinsky [FZ3] that cluster algebras are of finite type if and only if there is a seed $(A', \underline{x}')$ which is equivalent to $(A, \underline{u})$ such that the Cartan part $C_{A'}$ of A' is a Cartan matrix of finite type. The type of the Cartan matrix C_A is called the type of the cluster algebra $\mathcal{A}_A$.

For example if $C_{A'} = \begin{pmatrix} 2 & -1 & 0 & \cdots & 0 \\ -2 & 2 & -1 & \cdots & 0 \\ & & \cdots & \cdots & \\ 0 & 0 & \cdots & 2 & -1 \\ 0 & & \cdots & -1 & 2 \end{pmatrix}$, then the cluster algebra is called of type C_n .

Now we recall some basics on 2-CY triangulated categories.

Fix an algebraically closed field k . We say a triangulated category C is k -linear provided all Hom-spaces in C are k -spaces and the compositions of maps are k -linear. The k -linear triangulated categories in this paper will be assumed Hom-finite and Krull-Remak-Schmidt, i.e. $\dim_k \text{Hom}(X, Y) < \infty$ for any two objects X and Y in C , and every object decomposes into a finite direct sum of objects having local endomorphism rings. For using Hom-infinite triangulated categories to study cluster algebras, we refer the recent paper [Pla1&Pla2].

A k -linear triangulated category C is called 2-Calabi-Yau (2-CY for short) if there is a functorial isomorphism $\text{Hom}(X, Y) \cong D\text{Hom}(Y, X[2])$ for objects $X, Y \in C$, where $D = \text{Hom}_k(-, k)$ [K1].

Examples of 2-CY triangulated categories are the cluster categories of abelian hereditary categories with tilting objects [BMRRT, Ke1]; the Hom-finite generalized cluster categories of algebras with global dimension of at most 2 [Am]; the stable categories of stably 2-CY categories [BIRS], cluster tubes [BKL] and some others, please see the survey [K3].

Now we recall notions of cluster tilting objects, and of maximal rigid objects from [BIRS] [BMV].

Definition 2.1. Let T be an object of a 2-CY triangulated category C .

1. T is called rigid provided $\text{Ext}^1(T, T) = 0$.
2. T is called maximal rigid provided T is rigid and is maximal with respect to this property, i.e. if $\text{Ext}^1(T \oplus M, T \oplus M) = 0$, then $M \in \text{add}T$.
3. T is called cluster-tilting provided for any $M \in C$, if $\text{Ext}^1(M, T) = 0$, then $M \in \text{add}T$.

From the definition, any cluster tilting object is maximal rigid, but the converse is not true. It was proved in [BMV] that the cluster tube C_n has no cluster tilting objects, but maximal rigid objects. See [BIKR] for more examples. 2-CY triangulated categories with cluster tilting objects is important for the categorification of cluster algebras of skew-symmetric matrices, see the survey [K2] and the references there.

For a basic maximal rigid object $T = T_1 \oplus \cdots \oplus T_i \oplus \cdots \oplus T_n$ with all T_j indecomposable. For any $i \in [1, n]$, write $T = \bar{T} \oplus T_i$. Then there are two non-split triangles:

$$\begin{array}{ccccc} T_i^* & \xrightarrow{f_i} & E_i & \xrightarrow{g_i} & T_i \rightarrow T_i^*[1], \\ & & \downarrow f'_i & & \downarrow g'_i \\ T_i & \xrightarrow{f'_i} & E'_i & \xrightarrow{g'_i} & T_i^* \rightarrow T_i[1], \end{array}$$

where f_i and f'_i are minimal left $\bar{T}$ -approximations; g_i and g'_i are minimal right $\bar{T}$ -approximations. Furthermore T_i^* is indecomposable and $\bar{T} \oplus T_i^*$ is maximal rigid [IY BIRS]. Define $\mu_i(T) = \bar{T} \oplus T_i$, and call it the mutation of maximal rigid object T in direction i . It is easy to see that $\mu_i \circ \mu_i(T) = T$. The two triangles above are called exchange triangles.

Let $T = T_1 \oplus \cdots \oplus T_i \oplus \cdots \oplus T_n$ be a basic maximal rigid object in C . We define an integer matrix $A_T = (a_{ij})$ as follows:

$a_{ij} = \alpha_{ij} - \alpha'_{ij}$, where α_{ij} denotes the multiplicity of T_i as a direct summand of E'_j , α'_{ij} denotes the multiplicity of T_i as a direct summand of E_j . Note that $a_{ii} = 0$.

Our definition of the matrix A_T associated to T is the transpose of the matrix B_T associated to T defined in [BMV]. When the endomorphism algebra of T contains no loops or 2-cycles, then A_T, B_T are skew-symmetric matrices, and then $A_T = -B_T$. In general the matrices A_T, B_T are sign-skew-symmetric (see Lemma 1.2 in [BMV]).

We also note that

Remark 2.2. *We use our definition of the matrix A_T associated to T to replace the matrix B_T defined in [BMV] since we will prove the exchange formula for our cluster formulas of indecomposable rigid objects in C_n , and the cluster algebra realized by cluster tubes is of type C_{n-1} .*

Let C be a 2-CY triangulated category with maximal rigid objects. Suppose that for all maximal rigid objects $T = T_1 \oplus \cdots \oplus T_i \oplus \cdots \oplus T_n$, E_i and E'_i have no common direct summands for any $i \in [1, n]$. Then $\mu_i(A_T) = A_{\mu_i(T)}$ (equivalent to $\mu_i(B_T) = B_{\mu_i(T)}$, the latter was proved in [BMV]). In this case we say that the maximal rigid objects form a cluster structure in C as in [BMV].

In what follows, we will focus on cluster tubes, an special 2-CY triangulated category.

The tube $\mathcal{T}_n$ of rank n is the category of finite-dimensional nilpotent representations over k of the cyclic quiver $\vec{\Delta}_n$ with n vertices such that arrows are going from i to $i + 1$ (taken modulo n). It is a k -linear hereditary abelian category with Hom-finite, i.e. $\dim \text{Hom}(X, Y) < \infty$. Each indecomposable representation is uniserial, i.e. it has a unique composition series, and hence is determined by its socle and its length up to isomorphism. We denote by (a, b) in $\mathcal{T}_n$ the unique indecomposable object with socle $(a, 1)$ and quasi-length b , where $(a, 1)$ is the simple representation at vertex a , where $a \in [1, n]$ (see Fig.2 in Section 3). $\mathcal{T}_n$ has Auslander-Reiten sequences, and the Auslander-Reiten translation τ is an automorphism of $\mathcal{T}_n$, $\tau(a, b) = (a - 1, b)$. The cluster tube of rank n is defined in [BKL], see also [BMV], as the orbit category $C_n := D^b(\mathcal{T}_n)/\tau^{-1}[1]$, where $[1]$ is the shift functor of $D^b(\mathcal{T}_n)$. This category is a triangulated category such that the projection $\pi : D^b(\mathcal{T}_n) \rightarrow C_n$ is a triangle functor. It is also 2-CY [K].

C_n has Auslander-Reiten triangles which induced from ones in $D^b(\mathcal{T}_n)$. It is easy to see the indecomposable objects in $\mathcal{T}_n$ are also indecomposable in C_n (via the composite of the inclusion functor $\mathcal{T}_n \hookrightarrow D^b(\mathcal{T}_n)$ with projection $\pi : D^b(\mathcal{T}_n) \rightarrow C_n$) and all indecomposable objects in C_n are of this form. So we use the same (a, b) to denote the indecomposable object in C_n which is induced from the object (a, b) in $\mathcal{T}_n$.

By the definition of C_n , for two objects $X, Y \in \mathcal{T}_n$, $\text{Hom}_{C_n}(X, Y) \cong \text{Hom}_{\mathcal{T}_n}(X, Y) \amalg D\text{Hom}_{\mathcal{T}_n}(Y, \tau^2 X)$. As in [BMV], the maps from $\text{Hom}_{\mathcal{T}_n}(X, Y)$ are called $\mathcal{T}$ -maps and the maps from $D\text{Hom}_{\mathcal{T}_n}(Y, \tau^2 X)$ are called $\mathcal{D}$ -maps.

The indecomposable rigid objects are classified in [BMV]:

Proposition 2.3. *(a, b) is rigid if and only if $b \leq n - 1$.*

Denote by $T_i = (1, n - i)$, $i = 1, \dots, n - 1$. It is easy to see that $\bigoplus_{i=1}^{n-1} T_i$ is a maximal rigid object in C_n . We will use T to denote this maximal rigid object through the paper.

It was proved in [BMV] that the maximal rigid objects have no 2-cycles, and then form a cluster structure of C_n . Moreover there is a bijection between the indecomposable rigid objects and the cluster variables of the cluster algebra of type B_{n-1} .

For an object M in C_n , denote by $\text{add} M$ the subcategory of C_n consisting of (finite) direct sums of direct summands of M . For two subcategories $\mathcal{D}_1, \mathcal{D}_2$ of C_n , denote by $\mathcal{D}_1 * \mathcal{D}_2$ the full

subcategory of C_n consisting of object E such that there is a triangle $D_1 \rightarrow E \rightarrow D_2 \rightarrow D_1[1]$, where $D_i \in \mathcal{D}_i$, for $i = 1, 2$.

3 Index and the cluster map $X_?$

Let C_n be the cluster tube of rank n and $\mathcal{F} = \mathbf{Q}(x_1, x_2, \dots, x_{n-1})$ be the field of rational functions in indeterminates $x_1, x_2, \dots, x_{n-1}$. We always assume that $n \geq 2$ in the rest of the paper.

In this section, we will define the cluster map $X_?$ from the set of indecomposable rigid objects in C_n to $\mathcal{F}$ by using the geometric information of the indecomposable rigid objects. We will give an explicit expression of X_M as a Laurent polynomial of $x_1, \dots, x_{n-1}$ according to which subset M belongs to, where the set of indecomposable rigid objects in C_n is divided into three disjoint subsets.

Fix $T := \bigoplus_{i=1}^{n-1} T_i$, where $T_i = (1, n-i)$ is as before. T is a maximal rigid object in C_n . Let $K_0^{split}(T)$ be the split-Grothendieck group of $\text{add}T$, i.e. the free abelian group with a basis consisting of isomorphism classes of indecomposable direct summands of T . We use $[T']$ to denote the image in $K_0^{split}(T)$ of $T' \in \text{add}T$.

Let $\mathcal{D} = \text{add}T[-1] * \text{add}T$. For an object X of $\mathcal{D}[1]$, there exists a triangle

$$T_X'' \rightarrow T_X' \xrightarrow{f} X \rightarrow T_X''[1]$$

where $T_X'', T_X' \in \text{add}T$. It follows that f is a right $\text{add}T$ -approximation. We define the index $\text{ind}_T(X) = [T_X'] - [T_X''] \in K_0^{split}(T)$ as in [Pa1, DK, Pla]. It was proved in [ZZ] that any rigid object belongs to $\mathcal{D}[1]$ (also in $\mathcal{D}$). The next lemma tells us how to get the right $\text{add}T$ -approximation of any indecomposable rigid object in C_n .

Lemma 3.1. *For any indecomposable rigid object (a, b) in $\mathcal{T}_n$, every $\text{add}T$ -approximation of (a, b) in $\mathcal{T}_n$ is an $\text{add}T$ -approximation of (a, b) in C_n .*

Proof. It is easy to see that every $\mathcal{D}$ -map in $\text{Hom}_{C_n}(T_i, (a, b))$ factors through a $\mathcal{D}$ -map in $\text{Hom}_{C_n}(T_1, (a, b))$ (Lemma 2.2 in [BMV]). If there are no non-zero $\mathcal{D}$ -maps in $\text{Hom}_{C_n}(T, (a, b))$, then the $\text{add}T$ -approximation of (a, b) in $\mathcal{T}_n$ is an $\text{add}T$ -approximation of (a, b) in C_n . Now suppose there are non-zero $\mathcal{D}$ -maps in $\text{Hom}_{C_n}(T, (a, b))$, then $\dim \text{Hom}_{\mathcal{T}_n}((a, b), \tau^2 T_1) = 1$. So it is sufficient to show that for any nonzero $\mathcal{D}$ -map f in $\text{Hom}_{C_n}(T_1, T_1)$ and any nonzero $\mathcal{T}$ -map h in $\text{Hom}_{C_n}(T_1, (a, b))$ the composition hf is not zero. The triangle involving f is $(2, 2n-2) \rightarrow T_1 \xrightarrow{f} T_1 \xrightarrow{g} (1, 2n-2)$ where g is a $\mathcal{T}$ -map. We know that $\alpha g = 0$ for any $\mathcal{T}$ -map α in $\text{Hom}_{C_n}((1, 2n-2), (a, b))$, so h can not factor through g , and then $hf \neq 0$. Thus we have the assertion. $\square$

Theorem 3.2. *For any indecomposable rigid object (a, b) in C_n ,*

$$\text{ind}_T(a, b) = \begin{cases} [T_1] - [T_{n-a+1}] - [T_{2n-a-b}] & \text{if } a + b \geq n + 1 \\ [T_{n-a-b+1}] - [T_{n-a+1}] & \text{if } a + b \leq n. \end{cases}$$

Proof. For $a = n$, there is a triangle $(1, b) \rightarrow 0 \rightarrow (n, b) \rightarrow (1, b)[1]$, so $\text{ind}_T(n, b) = -[(1, b)]$. For $1 \leq a \leq n-1$ and $a+b \geq n+1$, there is a minimal right $\text{add}T$ -approximation $f : T_1 \rightarrow (a, b)$ in $\mathcal{T}_n$. Then $\ker(f) = (1, a-1)$ and $\text{coker}(\tau^{-1}f) = (1, a+b-n)$. We get a triangle

$$C \rightarrow T_1 \xrightarrow{f} (a, b) \rightarrow C[1]$$

in C_n and an exact sequence

$$0 \rightarrow \text{coker}(\tau^{-1}f) \rightarrow C \rightarrow \ker(f) \rightarrow 0$$

in $\mathcal{T}_n$ (cf. [Y]). Since $\text{Ext}_{\mathcal{T}_n}^1((1, a-1), (1, a+b-n)) = 0$, then $C \cong (1, a-1) \oplus (1, a+b-n)$. Hence $\text{ind}_T(a, b) = [T_1] - [T_{n-a+1}] - [T_{2n-a-b}]$.

For $a+b \leq n$, there is a minimal right $\text{add}T$ -approximation $f : T_{n-a-b+1} \rightarrow (a, b)$ in $\mathcal{T}_n$. Then $\ker(f) = (1, a-1)$ and $\text{coker}(\tau^{-1}f) = 0$. Hence $\text{ind}_T(a, b) = [T_{n-a-b+1}] - [T_{n-a+1}]$. $\square$

Let $B = \text{End}_{C_n}(T[-1])$, then $F := \text{Hom}(T[-1], -) : \mathcal{D}/\text{add}T \rightarrow \text{mod } B$ is an equivalence of categories.

Definition 3.3. For any indecomposable rigid object M in C_n , we define

$$X_M = x^{\text{ind}_T M} \sum_{e \in \mathbb{N}^{n-1}} \chi(\text{Gr}_e(\text{Hom}(T[-1], M))) x^{-\iota(e)}$$

where the sum takes over all dimension vectors e such that there exist an object Y in $\mathcal{D} \cap \mathcal{D}[1]$ with $e = \underline{\dim} \text{Hom}_{C_n}(T[-1], Y)$; where $\iota(e) = \text{ind}_T Y + \text{ind}_T Y[1]$ and χ is the Euler-Poincaré characteristic of $\text{Gr}_e(\text{Hom}(T[-1], M))$ ([K2]).

For any rigid object $N = \bigoplus_{i=1}^m M_i$ in C_n with M_i indecomposable, we define

$$X_N = \prod_{i=1}^m X_{M_i}.$$

In [Pla1][Pla2], similar definition is given respect to a fixed rigid object T . But his definition needs additional assumption that the simple B -module at each vertex can be lifted to an object in $\mathcal{D} \cap \mathcal{D}[1]$ through the functor F and then any finite-dimensional B -module can be lifted through F to an object in $\mathcal{D} \cap \mathcal{D}[1]$. In our situation, the assumption is not true and we omit the B -modules which can not be lifted through F to $\mathcal{D} \cap \mathcal{D}[1]$ in the definition above. In fact simple B -module S_1 corresponding to T_1 can not lift to any object in $\mathcal{D} \cap \mathcal{D}[1]$ by functor F (see the dimension formula stated before Lemma 3.5).

We give some remarks about the definition:

Remark 3.4. We will point out in Lemma 3.4 that for an indecomposable rigid object M , if a B -submodule of FM can not be lifted to $\mathcal{D} \cap \mathcal{D}[1]$, then so are other B -submodules of FM with the same dimension.

Remark 3.5. Why does not $\iota(e)$ depend on the choice of such an Y ? We will show this in the proof of Theorem 3.7.

Following [V][Y], the set of indecomposable objects in $\mathcal{D}[1]$ is the set of indecomposable objects (a, b) satisfying either (1) (a, b) is rigid or (2) $n \leq b \leq 2n - 2$ and $a + b \leq 2n - 1$. We divide $\mathcal{D}$ into five subsets (see Fig.1):

$$\begin{aligned} O &= \{(a, b) \mid a = 1, b \leq n - 1\} \\ I &= \{(a, b) \mid 2 \leq a \leq n - 1, a + b \leq n\} \\ II &= \{(a, b) \mid a + b \geq n + 1, b \leq n - 1\} \\ III &= \{(a, b) \mid a + b \leq 2n - 1, a \neq 1, b \geq n\} \\ IV &= \{(a, b) \mid a + b = 2n, a \neq 1, b \geq n\} \end{aligned}$$




Fig.1 Five disjoint subsets of $\mathcal{D}$.

It is easy to see that for any indecomposable object (a, b) in $\mathcal{D}$, we have

$$\dim \text{Hom}(T_1[-1], (a, b)) = \begin{cases} 0 & \text{if } (a, b) \in O \text{ or } I; \\ 2 & \text{if } (a, b) \in II \text{ or } III; \\ 1 & \text{if } (a, b) \in IV. \end{cases}$$

So we have the following fact.

Lemma 3.6. *For any indecomposable rigid object M in C_n , if Y is an object in $\mathcal{D}$ such that FY is a B -submodule of FM , then $Y \in \mathcal{D} \cap \mathcal{D}[1]$ if and only if $\dim \text{Hom}(T_1[-1], Y) \neq 1$.*

Proof. We know that for object Y in $\mathcal{D}$, $Y \in \mathcal{D} \cap \mathcal{D}[1]$ if and only if all indecomposable summands of Y are in $O \cup I \cup II \cup III$.

Since $\dim \text{Hom}(T_1[-1], M) \leq 2$, so $\dim \text{Hom}(T_1[-1], Y) = 2$ implies that $\dim \text{Hom}(T_1[-1], M) = 2$ and . Then there is a $\mathcal{T}$ -map f in $\text{Hom}(T_1[-1], M)$ such that f is a generator of B -module FM by the proof of Theorem 3.2. So $FY \cong FM$ and then $Y \cong M \oplus T'$, where $T' \in \text{add} T$. Hence $Y \in \mathcal{D} \cap \mathcal{D}[1]$.

When $\dim \text{Hom}(T_1[-1], Y) = 1$, Y has some summand in IV , then $Y \notin \mathcal{D} \cap \mathcal{D}[1]$.

When $\dim \text{Hom}(T_1[-1], Y) = 0$, all summands of Y are in O or I , then $Y \in \mathcal{D} \cap \mathcal{D}[1]$.

Thus the statement holds. □

Let $E_i : \text{Hom}(T_i[-1], Y) \rightarrow \text{Hom}(T[-1], Y)$ be the natural embedding and e_i be the dimension vector of S_i where S_i is the simple module of B corresponding to T_i . In the following expression of x_m , when $m = n$, then x_n is taken to 1.

Theorem 3.7. The X_M in Definition 3.3 is well-defined and for any indecomposable rigid object (a, b) ,

$$X_{(a,b)} = \begin{cases} x_{n-b} & \text{if } (a, b) \in O; \\ \sum_{k=0}^b \frac{x_{n-a-b+1}x_{n-a+2}}{x_{n-a-k+1}x_{n-a-k+2}} & \text{if } (a, b) \in I; \\ x_1x_{n-a+2}x_{2n-a-b+1} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{x_{n-a-k+1}x_{n-a-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-1} \frac{1}{x_{2n-a-b-l}x_{2n-a-b-l+1}} \right) \right) & \text{if } (a, b) \in II. \end{cases}$$

Proof. For the first statement, we have only to show that if two objects $Y', Y'' \in \mathcal{D}$ with $\underline{\dim}FY' = \underline{\dim}FY''$ and an indecomposable rigid object $X \in \mathcal{D}$ satisfy both FY' and FY'' are B -submodule of FX , then $\text{ind}_T Y' + \text{ind}_T Y'[1] = \text{ind}_T Y'' + \text{ind}_T Y''[1]$. We will show this in the following detailed calculations. In the following analysis, if $FY \cong FY_1$ as a B -submodule of FM , then $Y \oplus T' \cong Y_1 \oplus T''$, for some $T', T'' \in \text{add}T$. Hence we have $\text{ind}_T(Y \oplus T') + \text{ind}_T(Y \oplus T')[1] = \text{ind}_T(Y_1 \oplus T'') + \text{ind}_T(Y_1 \oplus T'')[1]$. Then $\text{ind}_T Y + \text{ind}_T Y[1] = \text{ind}_T Y_1 + \text{ind}_T Y_1[1]$. So we will consider only the situation of Y, Y_1 such that FY, FY_1 are not isomorphic to each other as submodules of FM .

1. The case of $(a, b) \in O$. $(a, b) \in \text{add}T$, hence we have $\text{ind}_T(a, b) + \text{ind}_T(a, b)[1] = 0$.
2. The case of $(a, b) \in I$. It is easy to compute the dimension vector of $F((a, b))$:

$$\dim \text{Hom}(T_i[-1], (a, b)) = \begin{cases} 1 & \text{if } n-a-b+2 \leq i \leq n-a+1 \\ 0 & \text{otherwise.} \end{cases}$$

For any nonzero element f in $\text{Hom}(T_i[-1], (a, b))$, $n-a-b+2 \leq i \leq n-a+1$, f is a $\mathcal{T}$ -map. Then $E_i(f)B = \oplus_{j=i}^{n-a+1} E_j(\text{Hom}(T_j[-1], (a, b)))$. Then $F(Y)$ is a B -submodule of $F((a, b))$ with $\dim \text{Hom}(T_1[-1], Y) \neq 1$ if and only if $F(Y) \cong E_i(f)B$ for some $f \in \text{Hom}(T_i[-1], (a, b))$, $n-a-b+2 \leq i \leq n-a+1$. Hence $FY \cong F(a, b')$, $0 \leq b' \leq b$. Then $\text{ind}_T Y + \text{ind}_T Y[1] = \text{ind}_T(a, b') + \text{ind}_T(a, b')[1]$. Moreover, for $0 \leq b' \leq b$, $\chi(\text{Gr}_{\underline{\dim}(F(a, b'))}(\text{Hom}(T[-1], (a, b)))) = 1$. Hence, by Definition 3.3, we have that $X_{(a,b)} = \sum_{k=0}^b \frac{x_{n-a-b+1}x_{n-a+2}}{x_{n-a-k+1}x_{n-a-k+2}}$.

3. The case of $(a, b) \in II$.

(a) $b = n-1$. In this case,

$$\dim \text{Hom}(T_i[-1], (a, b)) = \begin{cases} 2 & \text{if } 1 \leq i \leq n-a+1 \\ 0 & \text{otherwise.} \end{cases}$$

There is a basis $\{E_1(f_1), E_1(g_1), \dots, E_{n-a+1}(f_{n-a+1}), E_{n-a+1}(g_{n-a+1})\}$ of $F((a, b))$ where f_i is a $\mathcal{T}$ -map in $\text{Hom}(T_i[-1], (a, b))$ and g_i is a $\mathcal{D}$ -map in $\text{Hom}(T_i[-1], (a, b))$. Then $E_i(f_i)B = \oplus_{j=i}^{n-a+1} E_j(f_j)$ ($2 \leq i \leq n-a+1$), $E_i(g_i)B = \oplus_{j=i}^{n-a+1} E_j(g_j)$ ($1 \leq i \leq n-a+1$) and $E_1(f_1)B = F((a, b))$. So if $F(Y)$ is a B -submodule of $F((a, b))$ with $\dim \text{Hom}(T_1[-1], Y) \neq 1$, there are five cases:

- i. $Y \cong (a, b)$ with $\underline{\dim}(FY) = \sum_{k=1}^{n-a+1} 2e_k$. In this case, $\chi = 1$.
- ii. $F(Y) \cong E_i(\text{Hom}(T_i, (a, b)))B$ with $\underline{\dim}(FY) = \sum_{k=i}^{n-a+1} 2e_k$, $2 \leq i \leq n-a+1$ (when $a \leq n-1$). Then $Y \cong (a, b') \oplus (a, b')$, $1 \leq b' \leq n-a$. In this case, $\chi = 1$.

- iii. $F(Y) \cong E_i(f_i)B \oplus E_i(g_j)B$ with $\underline{\dim}(FY) = \sum_{k=i}^{n-a+1} e_k + \sum_{l=j}^{n-a+1} e_l$, $2 \leq i < j \leq n-a+1$ (when $a \leq n-2$). Then $Y \cong (a, b') \oplus (a, b'')$, $1 \leq b'' < b' \leq n-a$. In this case, $\chi = 2$.
- iv. $F(Y) \cong E_i(f_i)B$ with $\underline{\dim}(FY) = \sum_{k=i}^{n-a+1} e_k$, $2 \leq i \leq n-a+1$ (when $a \leq n-1$). Then $Y \cong (a, b')$, $1 \leq b' \leq n-a$. In this case, $\chi = 2$.
- v. $Y \cong 0$ with $\underline{\dim}(FY) = 0$. In this case, $\chi = 1$.

Hence, by Definition 3.3, we have that $X_{(a,n-1)} = x_1 x_{n-a+2}^2 \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{x_{n-a-k+1} x_{n-a-k+2}} \right)^2 \right)$.

(b) $b \leq n-2$. In this case,

$$\dim \text{Hom}(T_i[-1], (a, b)) = \begin{cases} 2 & \text{if } 1 \leq i \leq n-a+1 \\ 1 & \text{if } n+2-a \leq i \leq 2n-a-b \\ 0 & \text{otherwise.} \end{cases}$$

There is a basis $\{E_1(f_1), \dots, E_{n-a+1}(f_{n-a+1}), E_1(g_1), \dots, E_{2n-a-b}(g_{2n-a-b})\}$ of $F((a, b))$ where f_i is a $\mathcal{T}$ -map in $\text{Hom}(T_i[-1], (a, b))$ and g_i is a $\mathcal{D}$ -map in $\text{Hom}(T_i[-1], (a, b))$. Then $E_i(g_i)B = \oplus_{j=i}^{2n-a-b} E_j(g_j)$ ($1 \leq i \leq 2n-a-b$) and $E_1(f_1)B = F((a, b))$. So if $F(Y)$ is a B -submodule of $F((a, b))$ with $\dim \text{Hom}(T_1[-1], Y) \neq 1$, there are eight cases:

- i. $Y \cong (a, b)$ with $\underline{\dim}(FY) = \sum_{k=1}^{n-a+1} e_k + \sum_{l=1}^{2n-a-b} e_l$. In this case, $\chi = 1$.
- ii. $F(Y) \cong E_i(f_i)B$ with $\underline{\dim}(FY) = \sum_{k=i}^{n-a+1} e_k$, $2 \leq i \leq n-a+1$ (when $a \leq n-1$). Then $Y \cong (a, b')$, $1 \leq b' \leq n-a$. In this case, $\chi = 1$.
- iii. $F(Y) \cong E_i(g_i)B$ with $\underline{\dim}(FY) = \sum_{l=i}^{2n-a-b} e_l$, $n-a+2 \leq i \leq 2n-a-b$. Then $Y \cong (a+b-n+1, b'')$, $1 \leq b'' \leq n-b-1$. In this case, $\chi = 1$.
- iv. $F(Y) \cong E_i(g_i)B$ with $\underline{\dim}(FY) = \sum_{l=i}^{2n-a-b} e_l$, $2 \leq i \leq n-a+1$ or $F(Y) \cong BE_i(f_i) \oplus BE_{n-a+2}(g_{n-a+2})$ with $\underline{\dim}(FY) = \sum_{k=i}^{n-a+1} e_k + \sum_{l=n-a+2}^{2n-a-b} e_l$, $2 \leq i \leq n-a+1$ (when $a \leq n-1$). Then $Y \cong (a+b-n+1, b'')$, $n-b \leq b'' \leq 2n-a-b-1$ or $Y \cong (a, b') \oplus (a+b-n+1, n-b-1)$, $1 \leq b' \leq n-a$. Moreover, $\text{ind}_T((a+b-n+1, b'+n-b-1)) = \text{ind}_T(((a, b') \oplus (a+b-n+1, n-b-1)))$ and $\text{ind}_T((a+b-n, b'+n-b-1)) = \text{ind}_T(((a-1, b') \oplus (a+b-n, n-b-1)))$, $1 \leq b' \leq n-a$. In this case, $\chi = 2$.
- v. $F(Y) \cong E_i(f_i)B \oplus E_j(g_j)B$ with $\underline{\dim}(FY) = \sum_{k=i}^{n-a+1} e_k + \sum_{l=j}^{2n-a-b} e_l$, $2 \leq i \leq n-a+1$, $n-a+3 \leq j \leq 2n-a-b$ (when $a \leq n-1$ and $b \leq n-3$). Then $Y \cong (a, b') \oplus (a+b-n+1, b'')$, $1 \leq b' \leq n-a$, $1 \leq b'' \leq n-b-2$. In this case, $\chi = 1$.
- vi. $F(Y) \cong E_i(f_i)B \oplus E_i(g_i)B$ with $\underline{\dim}(FY) = \sum_{k=i}^{n-a+1} e_k + \sum_{l=i}^{2n-a-b} e_l$, $2 \leq i \leq n-a+1$ (when $a \leq n-1$). Then $Y \cong (a, b') \oplus (a+b-n+1, b'-b+n-1)$, $1 \leq b' \leq n-a$. In this case, $\chi = 1$.
- vii. $F(Y) \cong E_i(f_i)B \oplus E_j(g_j)B$ with $\underline{\dim}(FY) = \sum_{k=i}^{n-a+1} e_k + \sum_{l=j}^{2n-a-b} e_l$, $2 \leq i < j \leq n-a+1$ or $F(Y) \cong BE_i(f_i) \oplus BE_j(g_j)$ with $\underline{\dim}(FY) = \sum_{k=i}^{n-a+1} e_k + \sum_{l=j}^{2n-a-b} e_l$, $2 \leq j < i \leq n-a+1$ (when $a \leq n-2$). Then $Y \cong (a, b') \oplus (a+b-n+1, b'')$, $a+1 \leq a+b-n+1+b'' < a+b' \leq n$ or $Y \cong (a, b') \oplus (a+b-n+1, b'')$, $a+1 \leq a+b' < a+b-n+1+b'' \leq n$. Moreover, $\text{ind}_T((a, b') \oplus (a+b-n+1, b'')) = \text{ind}_T((a, b-n+b''+1) \oplus (a+b-n+1, n-b-b'-1))$ and

$$\text{ind}_T((a-1, b') \oplus (a+b-n, b'')) = \text{ind}_T((a-1, b-n+b''+1) \oplus (a+b-n, n-b-b'-1)).$$

In this case, $\chi = 2$.

viii. $Y \cong 0$ with $\underline{\dim}(FY) = 0$. In this case, $\chi = 1$.

Hence, by Definition 3.3, we have that

$$X_{(a,b)} = x_1 x_{n-a+2} x_{2n-a-b+1} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{x_{n-a-k+1} x_{n-a-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-1} \frac{1}{x_{2n-a-b-l} x_{2n-a-b-l+1}} \right) \right).$$

Thus the proof is completed. $\square$

Example 1. In C_3 , there are six indecomposable rigid objects: $T_1 = (1, 2)$, $T_2 = (1, 1)$, $\tau^{-1}T_1 = (2, 2)$, $\tau^{-1}T_2 = (2, 1)$, $\tau T_1 = (3, 2)$ and $\tau T_2 = (3, 1)$ (see Fig.2).




Fig.2 The tube of rank 3.

By Theorem 3.7, we have the following result where we reserve the details of the calculation:

$$X_{T_1} = x_1,$$

$$X_{T_2} = x_2,$$

$$X_{\tau^{-1}T_1} = \frac{x_1}{x_2} \left(\frac{x_2^2}{x_1^2} + \frac{1}{x_1^2} + 2\frac{1}{x_1} + 1 \right) = \frac{(x_1+1)^2 + x_2^2}{x_1 x_2^2},$$

$$X_{\tau^{-1}T_2} = \frac{1}{x_2} (x_1 + 1) = \frac{x_1+1}{x_2},$$

$$X_{\tau T_1} = \frac{1}{x_1} (x_2^2 + 1) = \frac{x_2^2+1}{x_1},$$

$$X_{\tau T_2} = \frac{1}{x_2} \left(\frac{x_2^2}{x_1} + \frac{1}{x_1} + 1 \right) = \frac{x_1+1+x_2^2}{x_1 x_2}.$$

4 Mutation relations

By Proposition 2.6 in [BMV], every maximal rigid object in C_n is in some wing of $(a, n-1)$ and there is a natural bijection between the set of maximal rigid object in the wing of $(a, n-1)$ and the set of tilting modules over path algebra $k\tilde{A}_{n-1}$ of linear quiver of type A_{n-1} . Hence using the complete description of all tilting modules of quivers of type A in [HR], one can easily to check that the following proposition for maximal rigid objects in C_n . In the following, for the pair of numbers of object in C_n , we always view $(a, 0)$ as zero object.

Proposition 4.1. *Let R be a maximal rigid object in C_n and (a, b) be an indecomposable summand of R . Then there are $(a, h - 1)$ and $(a + h, b - h)$ in $\text{add}R$ for some h , $1 \leq h \leq b$ and every indecomposable summand of R in the wing of (a, b) which is not isomorphic to (a, b) is in the wing of $(a, h - 1)$ or in the wing of $(a + h, b - h)$. When $b \neq n - 1$, there exists an i , $1 \leq i \leq n - b - 1$, such that either $(a, b + i)$ or $(a - i, b + i)$ is in $\text{add}R$.*

With this proposition and Theorem 1.1 in [BMV], we can give all exchange triangles in C_n .

Lemma 4.2. *Given two basic maximal rigid objects $T' \oplus \bar{T}$ and $T'' \oplus \bar{T}$ in C_n such that both T' and T'' are indecomposable. Then $\dim \text{Ext}_{C_n}^1(T', T'') = 1$ or 2 .*

Moreover, if $\dim \text{Ext}_{C_n}^1(T', T) = 2$, then the exchange triangles have the following form:

$$(a, n - 1) \rightarrow (a + h, n - h - 1) \oplus (a + h, n - h - 1) \rightarrow (a + h, n - 1) \rightarrow (a, n - 1)[1],$$

$$(a + h, n - 1) \rightarrow (a, h - 1) \oplus (a, h - 1) \rightarrow (a, n - 1) \rightarrow (a + h, n - 1)[1]$$

where $1 \leq a \leq n$, $1 \leq h \leq n - 1$.

If $\dim \text{Ext}_{C_n}^1(T', T) = 1$, then the exchange triangles have the following form:

$$(a, b) \rightarrow (a, b + i) \oplus (a + h, b - h) \rightarrow (a + h, b - h + i) \rightarrow (a, b)[1],$$

$$(a + h, b - h + i) \rightarrow (a + b + 1, i - 1) \oplus (a, h - 1) \rightarrow (a, b) \rightarrow (a + h, b - h + i)[1]$$

where $1 \leq a \leq n$, $b \leq n - 2$, $1 \leq h \leq b$, $1 \leq i \leq n - b - 1$.

Proof. Let $(a, n - 1) \oplus \bar{R}$ be a basic maximal rigid object, $1 \leq a \leq n$. Then there is an h , $1 \leq h \leq n - 1$, such that $(a, h - 1)$ and $(a + h, n - h - 1)$ are in $\text{add}((a, n - 1) \oplus \bar{R})$ and other indecomposable summands of $\bar{R}$ are in the wing of $(a, h - 1)$ or in the wing of $(a + h, n - h - 1)$. Now $((a + h, n - 1)) \oplus \bar{R}$ is a basic maximal rigid object. There is no nonzero map from the wing of $(a + h, n - h - 1)$ to $(a, n - 1)$ and any maps from the wing of $(a, h - 1)$ to $(a, n - 1)$ factor from $(a, h - 1)$. Since $\dim \text{Hom}_C((a, h - 1), (a, n - 1)) = 2$ and $\text{Hom}_C((a, h - 1), (a, n - 1))$ is not a cyclic $\text{End}_C((a, h - 1))$ -module, then the minimal right $\text{add} \bar{R}$ -approximation of $(a, n - 1)$ is $(a, h - 1) \oplus (a, h - 1) \rightarrow (a, n - 1)$. Similarly, we can get the minimal left $\text{add} \bar{R}$ -approximation of $(a, n - 1)$: $(a, n - 1) \rightarrow (a + h, n - h - 1) \oplus (a + h, n - h - 1)$. By Theorem 1.1 in [BMV], we have that the exchange triangles. In this case, $\dim \text{Ext}_{C_n}^1((a, n - 1), (a + h, n - 1)) = 2$, $1 \leq h \leq n - 1$.

Let $(a, b) \oplus \bar{R}$, $b \neq n - 1$, be a basic maximal rigid object, $1 \leq a \leq n$, $b \leq n - 2$. Then there is an i , $1 \leq i \leq n - b - 1$, such that either $(a, b + i)$ or $(a - i, b + i)$ is in $\text{add}((a, b) \oplus \bar{R})$. Assume that $(a, b + i)$ the "upper right" to (a, b) is in $\text{add}(\bar{R})$ and there is no j , $1 \leq j < i$, such that $(a, b + j) \in \text{add}(\bar{R})$. In the wing of (a, b) , there are $(a + h, b - h)$ and $(a, h - 1)$ in $\text{add}(\bar{R})$ for some h , $1 \leq h \leq b$. Now $((a + h, b - h + i)) \oplus \bar{R}$ is a basic maximal rigid object. Since $(a, b) \rightarrow (a, b + i) \oplus (a + h, b - h) \rightarrow (a + h, b - h + i) \rightarrow (a, b)[1]$ and $(a + h, b - h + i) \rightarrow (a + b + 1, i - 1) \oplus (a, h - 1) \rightarrow (a, b) \rightarrow (a + b + 1, i - 1)[1]$ are non-split triangles in $D^b(\mathcal{T}_n)$ and $\dim \text{Ext}_{C_n}^1((a, b), (a + h, b - h + i)) = 1$, so these two triangle are the exchange triangles between (a, b) and $(a + h, b - h + i)$. Note that $(a, b + i)$ is the "upper left" to $(a + h, b - h + i)$, so all exchange triangles are found. In this case, $\dim \text{Ext}_{C_n}^1((a, b), (a + h, b - h + i)) = 1$, $1 \leq i \leq n - b - 1$, $1 \leq h \leq b$. $\square$

The following is the main result in this section.

Theorem 4.3. Given two basic maximal rigid objects $T' \oplus \bar{T}$ and $T'' \oplus \bar{T}$ in C_n such that both T' and T'' are indecomposable. Then $X_{T'}X_{T''} = X_E + X_{E'}$, where $T' \rightarrow E \rightarrow T'' \rightarrow T'[1]$ and $T'' \rightarrow E' \rightarrow T' \rightarrow T''[1]$ are the exchange triangles.

Proof. 1. $\dim \text{Ext}_{C_n}^1(T', T) = 2$. By Lemma 4.2, we can assume that $T' = (a, n-1)$, $T'' = (a+h, n-1)$, $E = (a+h, n-h-1) \oplus (a+h, n-h-1)$ and $E' = (a, h-1) \oplus (a, h-1)$. We only need to consider two cases:

- (a) $a = 1$, $1 \leq h \leq n-1$. In this case, $(a, n-1), (a, h-1) \in O$, $(a+h, n-h-1) \in I$ and $(a+h, n-1) \in II$. Then

$$\begin{aligned} X_{(a, n-1)} &= x_1, \\ X_{(a+h, n-1)} &= x_1 x_{n-h+1} x_{n-h+1} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-h-1} \frac{1}{x_{n-h-k} x_{n-h-k+1}} \right)^2 \right), \\ X_{(a+h, n-h-1)} &= \sum_{k=0}^{n-h-1} \frac{x_1 x_{n-h+1}}{x_{n-h-k} x_{n-h-k+1}}, \\ X_{(a, h-1)} &= x_{n-h+1}. \end{aligned}$$

$$\text{Hence } X_{(a, n-1)} X_{(a+h, n-1)} = X_{(a, h-1)}^2 + X_{(a+h, n-h-1)}^2.$$

- (b) $2 \leq a \leq n-1$, $1 \leq h \leq n-a$. In this case, $(a, h-1) \in I$ and $(a, n-1), (a+h, n-1), (a+h, n-h-1) \in II$. Then

$$\begin{aligned} X_{(a, n-1)} &= x_1 x_{n-a+2} x_{n-a+2} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{x_{n-a-k+1} x_{n-a-k+2}} \right)^2 \right), \\ X_{(a+h, n-1)} &= x_1 x_{n-a-h+2} x_{n-a-h+2} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a-h} \frac{1}{x_{n-a-h-k+1} x_{n-a-h-k+2}} \right)^2 \right), \\ X_{(a+h, n-h-1)} &= x_1 x_{n-a-h+2} x_{n-a+2} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a-h} \frac{1}{x_{n-a-h-k+1} x_{n-a-h-k+2}} \right) \left(\sum_{l=0}^{n-a} \frac{1}{x_{n-a-l+1} x_{n-a-l+2}} \right) \right), \\ X_{(a, h-1)} &= \sum_{k=0}^{h-1} \frac{x_{n-a-h+2} x_{n-a+2}}{x_{n-a-k+1} x_{n-a-k+2}}. \end{aligned}$$

$$\text{Hence } X_{(a, n-1)} X_{(a+h, n-1)} = X_{(a, h-1)}^2 + X_{(a+h, n-h-1)}^2.$$

2. $\dim \text{Ext}_{C_n}^1(T', T) = 1$. By Lemma 4.2, we can assume that $T' = (a, b)$, $T'' = (a+h, b-h+i)$, $E = (a, b+i) \oplus (a+h, b-h)$ and $E' = (a+b+1, i-1) \oplus (a, h-1)$.

- (a) $(a, b) \in O$. In this case, $(a, b), (a, b+i), (a, h-1) \in O$, $(a+h, b-h), (a+h, b-h+i)$

$i), (a+b+1, i-1) \in I$. Then

$$\begin{aligned}
X_{(a,b)} &= x_{n-b}, \\
X_{(a+h,b-h+i)} &= \sum_{k=0}^{b-h+i} \frac{x_{n-b-i} x_{n-h+1}}{x_{n-a-h-k+1} x_{n-a-h-k+2}}, \\
X_{(a,b+i)} &= x_{n-b-i}, \\
X_{(a+h,b-h)} &= \sum_{k=0}^{b-h} \frac{x_{n-b} x_{n-h+1}}{x_{n-a-h-k+1} x_{n-a-h-k+2}}, \\
X_{(a+b+1,i-1)} &= \sum_{k=0}^{i-1} \frac{x_{n-b-i} x_{n-b}}{x_{n-a-(b+1)-k+1} x_{n-a-(b+1)-k+2}}, \\
X_{(a,h-1)} &= x_{n-h+1}.
\end{aligned}$$

Hence $X_{(a,b)} X_{(a+h,b-h+i)} = X_{(a,b+i)} X_{(a+h,b-h)} + X_{(a+b+1,i-1)} X_{(a,h-1)}$.

(b) $(a, b) \in I$. There are three cases:

- i. $a+b < n$, $1 \leq i \leq n-a-b$. In this case, $(a, b), (a+h, b-h+i), (a, b+i), (a+h, b-h), (a+b+1, i-1), (a, h-1) \in I$. Then

$$\begin{aligned}
X_{(a,b)} &= \sum_{k=0}^b \frac{x_{n-a-b+1} x_{n-a+2}}{x_{n-a-k+1} x_{n-a-k+2}}, \\
X_{(a+h,b-h+i)} &= \sum_{k=0}^{b-h+i} \frac{x_{n-a-b-i+1} x_{n-a-h+2}}{x_{n-a-h-k+1} x_{n-a-h-k+2}}, \\
X_{(a,b+i)} &= \sum_{k=0}^{b+i} \frac{x_{n-a-b-i+1} x_{n-a+2}}{x_{n-a-k+1} x_{n-a-k+2}}, \\
X_{(a+h,b-h)} &= \sum_{k=0}^{b-h} \frac{x_{n-a-b+1} x_{n-a-h+2}}{x_{n-a-h-k+1} x_{n-a-h-k+2}}, \\
X_{(a+b+1,i-1)} &= \sum_{k=0}^{i-1} \frac{x_{n-a-b-i+1} x_{n-a-b+1}}{x_{n-a-(b+1)-k+1} x_{n-a-(b+1)-k+2}}, \\
X_{(a,h-1)} &= \sum_{k=0}^{h-1} \frac{x_{n-a-h+2} x_{n-a+2}}{x_{n-a-k+1} x_{n-a-k+2}}.
\end{aligned}$$

Hence $X_{(a,b)} X_{(a+h,b-h+i)} = X_{(a,b+i)} X_{(a+h,b-h)} + X_{(a+b+1,i-1)} X_{(a,h-1)}$.

- ii. $a+b < n$, $n-a-b+1 \leq i \leq n-b-1$. In this case, $(a, b), (a, b+i), (a+h, b-h) \in I$

and $(a, b+i), (a+h, b-h+i), (a+b+1, i-1) \in II$. Then

$$\begin{aligned}
X_{(a,b)} &= \sum_{k=0}^b \frac{X_{n-a-b+1} X_{n-a+2}}{X_{n-a-k+1} X_{n-a-k+2}}, \\
X_{(a+h,b-h+i)} &= X_1 X_{n-a-h+2} X_{2n-a-b-i+1} \left(\frac{1}{X_1^2} + \left(\sum_{k=0}^{n-a-h} \frac{1}{X_{n-a-h-k+1} X_{n-a-h-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-i-1} \frac{1}{X_{2n-a-b-i-l} X_{2n-a-b-i-l+1}} \right) \right), \\
X_{(a,b+i)} &= X_1 X_{n-a+2} X_{2n-a-b-i+1} \left(\frac{1}{X_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{X_{n-a-k+1} X_{n-a-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-i-1} \frac{1}{X_{2n-a-b-i-l} X_{2n-a-b-i-l+1}} \right) \right), \\
X_{(a+h,b-h)} &= \sum_{k=0}^{b-h} \frac{X_{n-a-b+1} X_{n-a-h+2}}{X_{n-a-h-k+1} X_{n-a-h-k+2}}, \\
X_{(a+b+1,i-1)} &= X_1 X_{n-a-b+1} X_{2n-a-b-i+1} \left(\frac{1}{X_1^2} + \left(\sum_{k=0}^{n-a-b-1} \frac{1}{X_{n-a-b-k} X_{n-a-b-k+1}} \right) \left(\sum_{l=0}^{2n-a-b-i-1} \frac{1}{X_{2n-a-b-i-l} X_{2n-a-b-i-l+1}} \right) \right), \\
X_{(a,h-1)} &= \sum_{k=0}^{h-1} \frac{X_{n-a-h+2} X_{n-a+2}}{X_{n-a-k+1} X_{n-a-k+2}}.
\end{aligned}$$

Hence $X_{(a,b)} X_{(a+h,b-h+i)} = X_{(a,b+i)} X_{(a+h,b-h)} + X_{(a+b+1,i-1)} X_{(a,h-1)}$.

iii. $a+b=n$, $1 \leq i \leq n-b-1$. In this case, $(a+b+1, i-1) \in O$, (a, b) , $(a, b+i)$, $(a+h, b-h) \in I$ and $(a, b+i), (a+h, b-h+i) \in II$. Then

$$\begin{aligned}
X_{(a,b)} &= \sum_{k=0}^b \frac{X_1 X_{n-a+2}}{X_{n-a-k+1} X_{n-a-k+2}}, \\
X_{(a+h,b-h+i)} &= X_1 X_{n-a-h+2} X_{n-i+1} \left(\frac{1}{X_1^2} + \left(\sum_{k=0}^{n-a-h} \frac{1}{X_{n-a-h-k+1} X_{n-a-h-k+2}} \right) \left(\sum_{l=0}^{n-i-1} \frac{1}{X_{n-i-l} X_{n-i-l+1}} \right) \right), \\
X_{(a,b+i)} &= X_1 X_{n-a+2} X_{n-i+1} \left(\frac{1}{X_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{X_{n-a-k+1} X_{n-a-k+2}} \right) \left(\sum_{l=0}^{n-i-1} \frac{1}{X_{n-i-l} X_{n-i-l+1}} \right) \right), \\
X_{(a+h,b-h)} &= \sum_{k=0}^{b-h} \frac{X_{n-a-b+1} X_{n-a-h+2}}{X_{n-a-h-k+1} X_{n-a-h-k+2}}, \\
X_{(a+b+1,i-1)} &= X_{n-i+1}, \\
X_{(a,h-1)} &= \sum_{k=0}^{h-1} \frac{X_{n-a-h+2} X_{n-a+2}}{X_{n-a-k+1} X_{n-a-k+2}}.
\end{aligned}$$

Hence $X_{(a,b)} X_{(a+h,b-h+i)} = X_{(a,b+i)} X_{(a+h,b-h)} + X_{(a+b+1,i-1)} X_{(a,h-1)}$.

(c) $(a, b) \in II$. There are three cases:

i. $1 \leq h \leq n-a$. In this case, $(a, h-1), (a+b+1, i-1) \in I$ and $(a, b), (a, b+i), (a+$

$h, b-h), (a+h, b-h+1) \in II$. Then

$$\begin{aligned}
X_{(a,b)} &= x_1 x_{n-a+2} x_{2n-a-b+1} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{x_{n-a-k+1} x_{n-a-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-1} \frac{1}{x_{2n-a-b-l} x_{2n-a-b-l+1}} \right) \right), \\
X_{(a+h,b-h+i)} &= x_1 x_{n-a-h+2} x_{2n-a-b-i+1} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a-h} \frac{1}{x_{n-a-h-k+1} x_{n-a-h-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-i-1} \frac{1}{x_{2n-a-b-i-l} x_{2n-a-b-i-l+1}} \right) \right), \\
X_{(a,b+i)} &= x_1 x_{n-a+2} x_{2n-a-b-i+1} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{x_{n-a-k+1} x_{n-a-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-i-1} \frac{1}{x_{2n-a-b-i-l} x_{2n-a-b-i-l+1}} \right) \right), \\
X_{(a+h,b-h)} &= x_1 x_{n-a-h+2} x_{2n-a-b+1} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a-h} \frac{1}{x_{n-a-h-k+1} x_{n-a-h-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-1} \frac{1}{x_{2n-a-b-l} x_{2n-a-b-l+1}} \right) \right), \\
X_{(a+b+1,i-1)} &= \sum_{k=0}^{i-1} \frac{x_{n-a-b-i+1} x_{n-a-b+1}}{x_{n-a-(b+1)-k+1} x_{n-a-(b+1)-k+2}}, \\
X_{(a,h-1)} &= \sum_{k=0}^{h-1} \frac{x_{n-a-h+2} x_{n-a+2}}{x_{n-a-k+1} x_{n-a-k+2}}.
\end{aligned}$$

Hence $X_{(a,b)} X_{(a+h,b-h+i)} = X_{(a,b+i)} X_{(a+h,b-h)} + X_{(a+b+1,i-1)} X_{(a,h-1)}$.

- ii. $n-a+1 < h \leq b$. In this case, $(a+b+1, i-1), (a+h, b-h), (a+h, b-h+i) \in I$ and $(a, b), (a, b+i), (a, h-1) \in II$. Then

$$\begin{aligned}
X_{(a,b)} &= x_1 x_{n-a+2} x_{2n-a-b+1} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{x_{n-a-k+1} x_{n-a-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-1} \frac{1}{x_{2n-a-b-l} x_{2n-a-b-l+1}} \right) \right), \\
X_{(a+h,b-h+i)} &= \sum_{k=0}^{b-h+i} \frac{x_{n-a-b-i+1} x_{n-a-h+2}}{x_{n-a-h-k+1} x_{n-a-h-k+2}}, \\
X_{(a,b+i)} &= x_1 x_{n-a+2} x_{2n-a-b-i+1} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{x_{n-a-k+1} x_{n-a-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-i-1} \frac{1}{x_{2n-a-b-i-l} x_{2n-a-b-i-l+1}} \right) \right), \\
X_{(a+h,b-h)} &= \sum_{k=0}^{b-h} \frac{x_{n-a-b+1} x_{n-a-h+2}}{x_{n-a-h-k+1} x_{n-a-h-k+2}}, \\
X_{(a+b+1,i-1)} &= \sum_{k=0}^{i-1} \frac{x_{n-a-b-i+1} x_{n-a-b+1}}{x_{n-a-(b+1)-k+1} x_{n-a-(b+1)-k+2}}, \\
X_{(a,h-1)} &= x_1 x_{n-a+2} x_{2n-a-h+2} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{x_{n-a-k+1} x_{n-a-k+2}} \right) \left(\sum_{l=0}^{2n-a-h} \frac{1}{x_{2n-a-h-l+1} x_{2n-a-h-l+2}} \right) \right).
\end{aligned}$$

Hence $X_{(a,b)} X_{(a+h,b-h+i)} = X_{(a,b+i)} X_{(a+h,b-h)} + X_{(a+b+1,i-1)} X_{(a,h-1)}$.

- iii. $h = n-a+1$. In this case, $(a+h, b-h), (a+h, b-h+i) \in O, (a, h-1), (a+b+1, i-1) \in$

I and $(a, b), (a, b + i) \in II$. Then

$$\begin{aligned}
X_{(a,b)} &= x_1 x_{n-a+2} x_{2n-a-b+1} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{x_{n-a-k+1} x_{n-a-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-1} \frac{1}{x_{2n-a-b-l} x_{2n-a-b-l+1}} \right) \right), \\
X_{(a+h,b-h+i)} &= x_{2n-a-b-i+1}, \\
X_{(a,b+i)} &= x_1 x_{n-a+2} x_{2n-a-b-i+1} \left(\frac{1}{x_1^2} + \left(\sum_{k=0}^{n-a} \frac{1}{x_{n-a-k+1} x_{n-a-k+2}} \right) \left(\sum_{l=0}^{2n-a-b-i-1} \frac{1}{x_{2n-a-b-i-l} x_{2n-a-b-i-l+1}} \right) \right), \\
X_{(a+h,b-h)} &= x_{2n-a-b+1}, \\
X_{(a+b+1,i-1)} &= \sum_{k=0}^{i-1} \frac{x_{n-a-b-i+1} x_{n-a-b+1}}{x_{n-a-(b+1)-k+1} x_{n-a-(b+1)-k+2}}, \\
X_{(a,h-1)} &= \sum_{k=0}^{h-1} \frac{x_{n-a-h+2} x_{n-a+2}}{x_{n-a-k+1} x_{n-a-k+2}}.
\end{aligned}$$

$$\text{Hence } X_{(a,b)} X_{(a+h,b-h+i)} = X_{(a,b+i)} X_{(a+h,b-h)} + X_{(a+b+1,i-1)} X_{(a,h-1)}.$$

□

For the fixed maximal rigid object $T = \oplus_{i=1}^{n-1} T_i$, where $T_i = (1, n-i)$, from Lemma 4.2, we have

$$\text{that the matrix } A_T \text{ associated to } T \text{ is } A_T = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ -2 & 0 & 1 & \cdots & 0 \\ & & \cdots & \cdots & \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & & \cdots & -1 & 0 \end{pmatrix}. \text{ Its Cartan part is of type}$$

C_{n-1} .

Let $\text{indr}C_n$ be the set of indecomposable rigid objects in C_n , and χ_{A_T} be the set of cluster variables of skew-symmetrizable matrix A_T associated to T . Then we have the following:

Theorem 4.4. *The map $X_\gamma : \text{indr}C_n \longrightarrow \chi_{A_T} : M \mapsto X_M$ is a bijection. This bijection induces a bijection between the set of maximal rigid objects in C_n and the clusters of C_{n-1} . Furthermore the algebra generated by all X_M , where M runs through $\text{indr}C_n$, is isomorphic to any cluster algebra of type C_{n-1} .*

Proof. Recall that $T = \oplus_{i=1}^{n-1} T_i$. From the definition of X_M and Theorem 3.7, we have that $X_{T_i} = x_i$. Then X_γ sends the couple $(\{T_1, \dots, T_{n-1}\}, A_T)$ to the initial seed $(\{x_1, \dots, x_{n-1}\}, A_T)$ of the cluster algebra $\mathcal{A}_{A_T}$. It follows from Lemma 4.2 that the exchange graph of maximal rigid objects in C_n is connected. Then by Theorem 4.3, $X_\gamma(\text{indr}) \subset \chi_{A_T}$ and $X_\gamma : \text{indr}C_n \longrightarrow \chi_{A_T}$ is surjective. It follows from Theorem 3.7 that $X_\gamma : \text{indr}C_n \longrightarrow \chi_{A_T}$ is injective. Thus the first statement holds. The second statement also follows from the connection of the exchange graph of maximal rigid objects in C_n and that the initial maximal rigid object T goes to initial seed $(\{x_1, \dots, x_{n-1}\}, A_T)$ of the cluster algebra $\mathcal{A}_{A_T}$. The final statement follows from that the initial matrix A_T is of type C_{n-1} . □

Remark 4.5. *A bijection between from $\text{indr}C_n$ to the set of cluster variables of cluster algebras of type B_{n-1} was given in [BMV] by using the geometric description of cluster combinatorics of cluster algebras of type B_{n-1} given in [FZ2]. The cluster combinatorics of cluster algebras of*

type C_{n-1} is same as that of cluster algebras of type B_{n-1} [FZ2]. It follows that there is a same type bijection as in [BMV] between $\text{indr}C_n$ to the set of cluster variables of cluster algebras of type C_{n-1} . The theorem above gives an explicit expression of the bijection.

ACKNOWLEDGMENTS.

The authors would like to thank Dong Yang for his helpful discussion and suggestions.
The first-named author would like to thank Bin Li for his helpful discussion.

References

- [Am] C.Amiot. Cluster categories for algebras of global dimension 2 and quivers with potential. *Ann. Inst. Fourier* **59**, 2525-2590, 2009.
- [BKL1] M.Barot, D.Kussin and H.Lenzing. The Grothendieck group of a cluster category. *Jour. Pure and App. Algebra.* **212**, 33-46, 2008.
- [BKL2] M.Barot, D.Kussin and H.Lenzing. The cluster category of a canonical algebra. *Trans. Amer. Math. Soc.* **362**, no. 8, 4313–4330, 2010.
- [BIKR] I.Burban, O.Iyama, B.Keller and I.Reiten. Cluster tilting for one-dimensional hypersurface singularities. *Adv. Math.* **217**, 2443-2484, 2008.
- [BIRS] A.Buan, O.Iyama, I.Reiten and J.Scott. Cluster structure for 2–Calabi-Yau categories and unipotent groups. *Compo. Math.* **145**(4), 1035-1079, 2009.
- [BMV] A.Buan, R.Marsh and D.Vatne. Cluster structure from 2–Calabi-Yau categories with loops. *Mathematische Zeitschrift* **265**(4), 951-970, 2010.
- [BMRRT] A.Buan, R.Marsh, M.Reineke, I.Reiten and G.Todorov. Tilting theory and cluster combinatorics. *Advances in Math.* **204**, 572-618, 2006.
- [CC] P.Caldero and F.Chapoton. Cluster algebras as Hall algebras of quiver representations. *Comment. Math. Helv.* **81**, 595-616, 2006.
- [CCS] P.Caldero, F.Chapoton and R.Schiffler. Quivers with relations arising from clusters (A_n case). *Transactions of the AMS.* **358**, 1347-1364, 2006.
- [CK] P.Caldero and B.Keller. From triangulated categories to cluster algebras II. *Ann.Sci.Ecole.Norm.Sup.*(4) **39**, 983-1009,2006.
- [DK] R.Dehy and B.Keller. On the Combinatorics of Rigid Objects in 2-Calabi-Yau Categories. *International Mathematics Research Notices*, Vol.2008, Article ID run029, 17 pages.
- [Du] Dupont, G. An approach to non-simply laced cluster algebras. *J. Algebra* **320**, no. 4, 1626–1661, 2008.
- [DX] M.Ding and F.Xu The cluster character for cyclic quivers. *arXiv:1001.4360*.
- [FZ1] S.Fomin and A.Zelevinsky. Cluster Algebras I: Foundations. *Jour. Amer. Math. Soc.* **15**, no. 2, 497–529, 2002.

- [FZ2] S.Fomin and A.Zelevinsky. Y-system and generalized associahedra, *Ann of Math.* **158**, 977-1018, 2003.
- [FZ3] S.Fomin and A.Zelevinsky. Cluster algebras II: Finite type classification. *Invent. Math.* **154**, no.1, 63-121, 2003.
- [FZ4] S.Fomin and A.Zelevinsky. Cluster algebras: Notes for the CDM-03 Conference. CDM 2003: Current Developments in Mathematics. International Press, 2004.
- [FuKe] C.Fu and B.Keller. On cluster algebras with coefficients and 2-Calabi-Yau categories. *Trans.Amer.Math.Soc.* **362**, 859-895, 2010.
- [GLS1] C.Geiß, B.Leclerc and J.Shröer. Rigid modules over preprojective algebras. *Invent. math.* **165**(3), 589-632, 2006.
- [GLS2] C.Geiß, B.Leclerc and J.Shröer. Preprojective algebras and cluster algebras. *Trends in representation theory of algebras and related topics*, 253-283, EMS Ser.Congr.Rep. Eur.math.Soc. Zürich, 2008.
- [H] D.Happel. *Triangulated categories in the representation theory of quivers.* LMS Lecture Note Series, 119. Cambridge, 1988.
- [HR] D.Happel, C.M.Ringel. Construction of tilted algebras. *Representations of Algebras*, 125-144, 1981.
- [Hu] J. E. Humphreys. *Introduction to Lie Algebras and Representation Theory.* Springer-Verlag, New York, Heilelberg, Berlin, 1972.
- [I] O.Iyama. Higher dimensional Auslander-Reiten theory on maximal orthogonal subcategories. *Adv. Math.* **210**(2007), 22-50.
- [IY] O.Iyama and Y.Yoshino. Mutations in triangulated categories and rigid Cohen-Macaulay modules. *Invent. Math.* **172**, 117C168, 2008.
- [Ke1] B.Keller. Triangulated orbit categories. *Documenta Math.* **10**, 551-581, 2005.
- [Ke2] B.Keller. Cluster algebras, quiver representations and triangulated categories. *arXiv*: 0807.1960.
- [Ke3] B.Keller. Triangulated Calabi-Yau categories, *Trends in Representation Theory of Algebras (Zurich)* (A. Skowronski, ed.), European Mathematical Society, 2008, pp. 467C489.
- [KR] B.Keller and I.Reiten. Cluster-tilted algebras are Gorenstein and stably Calabi-Yau. *Adv. Math.* **211**, 123-151, 2007.
- [KZ] S.Koenig and B.Zhu. From triangulated categories to abelian categories— cluster tilting in a general framework. *Math. Zeit.* **258**, 143-160, 2008.
- [Pa1] Y.Palu. Cluster characters for 2-Calabi-Yau triangulated categories. *Ann.Inst.Fourier*, **58**, 133-171, 2008.

- [Pa2] Y.Palu. Grothendieck group and generalized mutation rule for 2–Calabi-Yau triangulated categories. *Jour.of Pure and Appl. Alg.* **213**, 1438-1449, 2009.
- [Pla1] P.Plamondon. Cluster characters for cluster categories with infinite-dimensional morphism spaces. *arXiv*. 1002.4956.
- [Pla2] P.Plamondon. Cluster algebras via cluster categories with infinite-dimensional morphism spaces. *arXiv*. 1004.0830.
- [Rin] C.M.Ringel. Some remarks concerning tilting modules and tilted algebras. Origin. Relevance. Future. An appendix to the Handbook of tilting theory, edited by Lidia Angeleri-Hügel, Dieter Happel and Henning Krause. Cambridge University Press (2007), LMS Lecture Notes Series 332.
- [V] D.Vatne. Endomorphism rings of maximal rigid objects in cluster tubes. *arXiv*: 0905.1796.
- [Y] D.Yang. Endomorphism algebras of maximal rigid objects in cluster tubes. *arXiv*: 1004.1303.
- [ZZ] Y.Zhou and B.Zhu. Maximal rigid subcategories in 2–Calabi-Yau triangulated categories. *arXiv*:1004.5475.